

Hawking radiation and thermodynamics of dynamical black holes in phantom dominated universe

Khiredine Nouicer*

Laboratory of Theoretical Physics (LPTh) and Department of Physics, Faculty of Sciences, University of Jijel, Bp 98 Ouled Aissa, Jijel 18000, Algeria

Abstract

The thermodynamic properties of dark energy-dominated universe in the presence of a black hole are investigated in the general case of a varying equation-of-state-parameter $w(a)$. We show that all the thermodynamics quantities are regular at the phantom divide crossing, and particularly the temperature and the entropy of the dark fluid are always positive definite. We also study the accretion process of a phantom fluid by black holes and the conditions required for the validity of the generalized second law of thermodynamics. As a results we obtain a strictly negative chemical potential and an equation-of-state parameter $w < -5/3$.

Keywords:GR black holes, dark energy theory, accretion of

1 Introduction

The discovery that the universe is currently undergoing a period of accelerating expansion, obtained from the observations of type Ia supernovae [1, 2], inaugurate an exciting era of intense theoretical research. A variety of possible solutions to understand the mechanism driving this accelerating expansion have been debated during this decade including the cosmological constant, exotic matter and energy, modified gravity, anthropic arguments, etc. The most favored ones are dark energy based models and modified gravity theories such $f(R)$ gravity [3], and Dvali-Gabadadze-Porrati model of gravity [4] with an equation of state (EoS) parameter $w = P/\rho < -1$ (where ρ and P are the energy density and pressure of the cosmic fluid, respectively). The dark energy is frequently modeled as an homogeneous scalar field, and according to the value of the EoS parameter w , three different cases can be distinguished: *quintessence scalar fields* ($-1 < w < -1/3$) with a positive kinetic term ; *cosmological constant* ($w = 1$), where only the potential term contributes to both the pressure and the energy density of the field; finally, scalar fields with a negative kinetic term dubbed

*E-mail:khnouicer@univ-jijel.dz

phantom fields ($w < -1$). In the latter case the universe will suffer a crucial fate where the energy density and the scale factor diverge in finite time, ripping apart all bound systems of the universe, before the universe approaches the so-called Big-Rip singularity [5, 6]. It is also well known that in phantom fluid models with big rip singularity, quantum gravity effects become dominant in the neighboring of the big rip time [7, 8, 9, 10]. However, even phantom fluid based models suffer from quantum instabilities [11] and violation of the strong and dominant classical energy conditions [12], the phantom fluids are favored by the cosmic microwave background experiments combined with large scale structure data, the Hubble parameter measurement and luminosity measurements of Type Ia supernovae [13].

An other important and growing field currently under investigation is related to the thermodynamic properties of an expanding universe [14]. Recent studies on phantom thermodynamics show that the entropy of the universe is negative [15], while the generalized second law of gravitational thermodynamics (GSL) is satisfied, $\dot{S}_f + \dot{S}_C \geq 0$, where S_f is the phantom fluid entropy and S_C is the entropy of the cosmological horizon [16]. An other point under debate is the influence of a non-zero chemical potential on the phantom thermodynamics and its relation with the GSL [17, 18].

In this paper the thermodynamic properties of black holes immersed in dark energy-dominated expanding universe and the accretion process of phantom energy onto black holes are investigated. The first paper dealing with the later process is due to Babichev et al [19], where ignoring the backreaction effect of the phantom fluid on the black hole, they found that the change rate of the black hole mass is negative. However, in recent scenarios where the backreaction is taken into account [20, 21], it is found that the mass of the black hole is always an increasing function in an expanding Friedman-Robertson-Walker (FRW) universe.

The organization of this paper is as follows: in Section 2 we review the exact solution recently obtained in [20, 21], and describing a black hole embedded in an expanding FRW universe. In section 3, we examine the Hawking radiation at the apparent horizon, and compare with magnitude of the phantom energy accretion process. We will show that the former is highly suppressed, particularly at late times. In section 4, we study in an unified and general way the thermodynamics of cosmological black hole embedded in an expanding (FRW) universe with a general EoS parameter $w(a)$, and particularly we obtain solutions realizing the crossing of the phantom divide line. In section 5, the stability of the solutions of section 4 under the quantum correction due to the conformal anomaly is established. In section 6, we study the conditions required for the validity of the GSL when the black hole is immersed in phantom energy-dominated FRW universe. Particularly, in order to protect the GSL, we obtain a critical mass of the black hole of the order of the solar mass for particular values of the parameter $\alpha_0 = -\mu_0 n_0 / \rho_0$, where μ_0 , n_0 , ρ_0 are the present day values of the chemical potential, the particle density and the energy density, respectively. Finally, we discuss and summarize our results in section 7.

2 Cosmological expanding black hole

The first solution of Einstein's theory of general relativity describing a black hole like object embedded in an expanding universe was introduced by McVittie in 1933 [22], and is given in isotropic coordinates by

$$ds^2 = -\frac{\left(1 - \frac{M_0}{2a(t)r}\right)^2}{\left(1 + \frac{M_0}{2a(t)r}\right)^2} dt^2 + a^2(t) \left(1 + \frac{M_0}{2a(t)r}\right)^4 (dr^2 + r^2 d\Omega^2), \quad (1)$$

where $a(t)$ is the scale factor and M_0 is the mass of the black hole in the static case. In fact, when $a(t) = 1$, it reduces to the Schwarzschild solution. When the mass parameter is zero, the McVittie reduces to a spatially flat FRW solution with the scale factor $a(t)$. The global structure of (1) has been studied and particularly it has been shown that the solution possesses a spacelike singularity on the 2-sphere $r = M_0/2$, and cannot describe an embedded black hole in an expanding spatially flat FLRW universe [23, 24]. On the other hand the McVittie solution is constrained by the non-accretion condition onto the central mass, and therefore is not suitable for a study of cosmic fluid accretion process onto black holes embedded in an expanding FRW universe.

In the following we adopt the new solution describing a black hole embedded in a spatially flat FRW universe [20, 21]

$$ds^2 = -\frac{B^2(r)}{A^2(r)} dt^2 + a^2(t) A^4(r) (dr^2 + r^2 d\Omega^2), \quad (2)$$

where $A(r) = \left(1 + \frac{GM_0}{2r}\right)$, $B(r) = \left(1 - \frac{GM_0}{2r}\right)$, $a(t)$ is the scale factor and M_0 is the mass of the black hole in the static case. In fact, when $a(t) = 1$, the solution (2) reduces to the Schwarzschild solution, while when the mass parameter is zero, it reduces to a spatially flat FRW solution with the scale factor $a(t)$.

Using the areal radius $\tilde{r} = r \left(1 + \frac{GM_0}{2r}\right)^2$ and $R = a\tilde{r}$, the metric takes the following suitable Painlevé-Gullstrand form

$$ds^2 = -\left[\left(1 - \frac{2GM_0 a}{R}\right) - \frac{R^2 H^2}{\left(1 - \frac{2GM_0 a}{R}\right)}\right] dt^2 + \left(1 - \frac{2GM_0 a}{R}\right)^{-1} dR^2 - 2RH \left(1 - \frac{2GM_0 a}{R}\right)^{-1} dt dR + R^2 d\Omega^2, \quad (3)$$

where $H = \dot{a}/a$ is the Hubble parameter and over dot stands for derivative with respect to the cosmic time. The term $R^2 H^2$ plays the role of variable cosmological constant. Now, we introduce the time transformation $t \rightarrow \bar{t}$ to remove the $dt dR$ term

$$d\bar{t} = F^{-1}(t, R) \left[dt + \frac{HR}{\left(1 - \frac{2GM_0a}{R}\right)^2 - H^2 R^2} dR \right], \quad (4)$$

where the integrating factor $F(t, R)$ satisfy

$$\partial_R F^{-1} = \partial_t \left[\frac{F^{-1}HR}{\left(1 - \frac{2GM_0a}{R}\right)^2 - H^2 R^2} \right]. \quad (5)$$

Substituting (4) into (3) and replacing $\bar{t} \rightarrow t$, we obtain

$$ds^2 = - \left[\left(1 - \frac{2GM_0a}{R}\right) - \frac{R^2 H^2}{\left(1 - \frac{2GM_0a}{R}\right)} \right] F^2 dt^2 \\ + \left[\left(1 - \frac{2GM_0a}{R}\right) - \frac{R^2 H^2}{\left(1 - \frac{2GM_0a}{R}\right)} \right]^{-1} dR^2 + R^2 d\Omega^2. \quad (6)$$

The apparent horizons (AH) are solutions of $h^{ab}\partial_a R \partial_b R = 0$, which leads to

$$\left(1 - \frac{2Gm_H(t)}{R} \mp RH\right)|_{R_A} = 0, \quad (7)$$

where we introduced the Hawking-Hayward quasi-local mass

$$m_H(t) = M_0 a(t). \quad (8)$$

A remarkable feature of this quantity is that it is coordinate-independent, and consequently is recognized as the physically relevant mass of the black hole. Obviously, it is always increasing in an expanding universe [21]. Therefore, the calculation of the change rate of the black hole mass will lead to opposite conclusions to that of Babichev et al. [19].

Discarding the unphysical branch with the lower sign in 7, the AH are given by

$$R_B = \frac{1}{2H} \left(1 - \sqrt{1 - 8G\dot{m}_H(t)}\right), \quad R_C = \frac{1}{2H} \left(1 + \sqrt{1 - 8G\dot{m}_H(t)}\right), \quad (9)$$

where R_C and R_B are the cosmological and the black hole AH, respectively. Note that the AH coincide at a time t_* defined by $\dot{a}(t_*) = 1/8GM_0$. This coincidence takes place in a future or past universe depending on the kind of fluid accretion onto the black hole.

Let us now consider that the fluid is described by a imperfect fluid with the stress-energy tensor given by

$$T_{\mu\nu} = (P + \rho) u_\mu u_\nu + P g_{\mu\nu} + q_\mu u_\nu + q_\nu u_\mu, \quad (10)$$

where $u^\mu = (\frac{A}{B}, 0, 0, 0)$ is the fluid four velocity and $q^\mu = (0, q, 0, 0)$ is a spatial vector field describing the radial energy current. Written in terms of the comoving AH, the solutions of the Einstein equations of motion are [21],

$$H = \frac{8\pi G}{3} R_A \rho, \quad 3H + \frac{2\dot{H}}{H} = -8\pi G R_A p. \quad (11)$$

Assuming a radial heat inflow ($q < 0$), the change rate of the black hole mass is then

$$\dot{m}_H = GaB^2 \mathcal{A} |q|, \quad (12)$$

where $\mathcal{A} = \int \int d\theta d\varphi \sqrt{g_\Sigma} = 4\pi r^2 a^2 A^4$. This relation shows that the Hawking-Hayward quasi-local mass is always increasing.

3 Hawking radiation of apparent horizon

For the apparent horizon to exist in an expanding universe we have the following condition

$$H(t) \leq \frac{1}{8Gm_H(t_*)}, \quad (13)$$

This condition requires us to consider massive objects for which $Gm_H H$ is a small quantity. Using the present day value of the Hubble parameter, we estimate the critical BH mass as $m_H(t_*) \lesssim 10^{23} M_\odot$. This is the condition for which the engulfing of the universe by the BH is prevented [25]. Using the areal radius $R = ar \left(1 + \frac{GM_0}{2r}\right)^2$, and solving for r , we obtain one physical solution given by

$$r_A = \frac{1}{2a} \left(R_A - Gm_H + \sqrt{R_A^2 - 2R_A Gm_H} \right). \quad (14)$$

For the solution to be real, we have to impose the condition

$$R_A > 2Gm_H, \quad (15)$$

which is true only in an expanding universe due to the relation $R_A(1 - HR_A) = 2Gm_H$. We note that the equality sign has been discarded in (15), since it leads to $HR_A = 0$, which does not hold for $m_H \neq 0$. The relation (15) means that we are dealing with systems which are small compared to the cosmological curvature, and that distances below $2l_{pl}$ are naturally excluded.

Assuming now the EoS, $p = w\rho$, and substituting (15) in Friedmann equations, one finds the following upper bounds on the densities

$$\rho(t) < \frac{3m_{pl}^6}{128\pi m_H^2(t)}, \quad |p| < \frac{3|w|m_{pl}^6}{128\pi m_H^2(t)}. \quad (16)$$

Therefore, in case of phantom energy driven expansion, the finite increase of densities with time, will avoid the big rip singularity. On the other, since the BH mass increases with time, the upper bounds (16), become very small. As an estimate of the energy density we have $\rho(t) < 0.02 \times 10^{-10} \text{ (GeV)}^4$, for the smallest super-massive BH detected in the dwarf Seyfert 1 galaxy POX 52 with $m_H \sim 10^5 M_\odot$ [26]. On the other hand if $m_H(t_*) \sim M_\odot$, we have $\rho(t) < 0.022 \text{ (GeV)}^4$. Even if we take m_H of the order of the Planck mass we have $\rho(t) < \frac{3m_{pl}^4}{128\pi}$.

We now consider the Hawking radiation from the cosmological AH. The temperature on the AH is defined by $T_A = |\kappa_A|/2\pi$, where κ_A is the surface gravity. In dynamical spacetime there is no timelike Killing vector, and the usual definition of the surface gravity may be modified. In this case the surface gravity is related to the so called trapping horizon. Here we follow the work of S. A. Hayward [27], where the surface gravity is defined by $K^b \nabla_{[a} K_{b]} = \kappa K_a$, ($a, b = 0, 1$), with $K^a = -\varepsilon^{ab} \nabla_b R$ the Kodama vector corresponding to the background given by Eq.(3). Evaluating all the quantities on the trapping horizon, we get the simplified form, $\kappa_A = \frac{1}{2} \square_h R$, where the metric h_{ab} is defined by $ds^2 = h_{ab} dx^a dx^b + R(x) d\Omega^2$. Performing the calculation, one finally finds

$$\kappa_A = \frac{m_H}{R_A^2} - H - \frac{\dot{H}}{2H}. \quad (17)$$

Writing m_H in terms of the AH, the temperature at the AH becomes

$$T_A = \frac{1}{4\pi R_A} \left| 1 - \frac{R_A}{H} (3H^2 + \dot{H}) \right|, \quad (18)$$

Expanding (18) to first order in Gm_H at the black hole AH and cosmic AH, we obtain

$$T_C = \frac{H}{2\pi} \left| 1 + \frac{\dot{H}}{2H^2} \right| - \frac{G\dot{m}_H}{2}, \quad (19)$$

$$T_B = \frac{1}{8\pi Gm_H} - \frac{GH^2 m_H}{2\pi} - \frac{4H^2 + \dot{H}}{4\pi H}, \quad (20)$$

respectively. Using (15) in the first factor, along with $R_C \leq 1/H$ in the second factor, and $\frac{\dot{H}}{H^2} = \frac{3}{2}(1+w)$, one obtains an instantaneous upper bound for temperature at cosmic AH

$$T_C < \frac{|1-3w|}{16\pi} \frac{m_{pl}^2}{m_H(t)}. \quad (21)$$

The relation (19) shows that the accretion process tends to lower the Hawking temperature at the cosmic AH. We note that in (20) we have not used the absolute value, as a consequence of the equivalence principle [28]. A similar behavior is also observed if one anticipates and use the scale factor (67) derived in section 4, for a phantom dominated era. In fact, one can see an unusual behavior of the temperature when approaching the time t_* . As it is shown in Fig. 1, the

temperature at the cosmic AH increases with time, reaches a maximum, then begins to decrease, and stops at the end time t_* . This means that the Hawking radiation at the cosmological AH will decrease rapidly in favor of a huge increasing of the accretion of phantom fluid. On the hand, the temperature of the black hole AH starts to fall at early stage because of the increasing of the black hole mass by accretion of phantom energy, and at late stage begins increasing. However, the latter strange behavior is mainly due to the absolute value in the definition of temperature.

An independent derivation of (16) and (21) can be performed by using the following simple arguments. Imposing positivity of the $(-g_{tt})$ component of the metric, and replacing the quasi-local mass by a density energy we have, $0 < -g_{tt} = 1 - 2Gm_H/R - \frac{R^2 H^2}{1 - 2Gm_H/R} = 1 - (8\pi G/3)\rho R^2 - \frac{R^2 H^2}{1 - (8\pi G/3)\rho R^2}$. Hence, $\rho < (1 - RH)/(8\pi GR^2/3)$. Using $R > 2Gm_H$, one obtains $\rho < 3m_{pl}^6/(32\pi m_H^2)$, which is consistent with (16). If one uses the Stefan-Boltzmann law $\rho = \sigma T^4$, and the quantum mechanical relation $R > 1/T$ [29], one finds $T < (3/4\pi\sigma)m_{pl}$, which of the same order of magnitude as (21) for $m_H = m_{pl}$. Finally, we point that the extremal case corresponding to $T_C = 0$ never occurs, since in this case we must have $R_C = \frac{1}{3H + \dot{H}/H^2}$, which contradicts the condition $R_C \geq 1/(2H)$.




Figure 1: Variation of temperature associated with the black hole and cosmic AH with time for $M_0 = 0.25$.

Now, neglecting the accretion of radiation in phantom energy dominated era, and taking into account only the semi-classical Hawking evaporation and the phantom energy accretion term, the differential equation for the black hole mass reads

$$\frac{dm_H}{dt} = -4\pi R_B^2 \sigma T_B^4 + m_H H, \quad (22)$$

where $\sigma = N\pi^2/120$ is the Stefan-Boltzmann constant for massless fields with effective degree of

freedom N . Substituting the Hawking temperature associated with the BH apparent horizon, we obtain

$$\frac{dm_H}{dt} = -\frac{\sigma H^2 \left| 1 - \frac{3}{4} (1 - \sqrt{1 - 8Gm_H H}) (1 - w) \right|^4}{16\pi^3 (1 - \sqrt{1 - 8Gm_H H})^2} + m_H H. \quad (23)$$

This is a complicated relation for m_H , whose behavior is shown in fig. 2. Since the two terms are in competition, there exist a transition time, the *phantom time*, after which the accretion process dominates and the BH mass increases. Consequently, the BH does not lose but gain mass due to high accretion of dark energy. Indeed, it is easy to show that the maximal rate gain mass for $m_H \neq 0$, is $(dm_H/dt)_{max} \sim m_{pl}^2$. It is important to note that the accretion term becomes predominant at earlier times for massive BH. Next, we perform the same analysis on the variation of the mass inside the cosmic AH. Let us define the ratio between the radiation and the accretion term

$$\eta_C(t) = \left| \frac{\dot{m}_{Haw}}{\dot{m}_{ph}} \right| = \frac{4\pi R_C^2 \sigma T_C^4}{m_H H}. \quad (24)$$

Substituting (18), and repeating the procedure leading to (21), one finds

$$\eta_C(t) < \frac{N |1 - 3w|^4}{245760\pi} \left(\frac{m_{pl}}{m_H} \right)^2. \quad (25)$$

This result clearly shows that the Hawking radiation at the cosmic AH is insignificant, even near the Planck scale.




Figure 2: Variation of total mass associated with the black hole AH (left panel) and cosmic AH (right panel) with time.

Finally, let us point out another crucial feature of the model of dynamical black hole considered in this paper. If the expansion of the universe is driven by phantom fluid, the black hole AH expands

while the cosmic AH shrinks as the universe expand until they meet at $R_{crit} = 1/(2H)$ at time t_* solution of (13). At times $t > t_*$, both the AH disappear living a proper singularity, well before the big-rip singularity is reached. The question if this singularity is naked or located inside the AH, and its connection with the violation of the Cosmic Censorship Conjecture (CCH), is still under debates [31, 32]. To avoid discussing this topics, which are out of the scope of the present paper, and the fact that the phantom driven expansion of the universe is non singular, and that the radiation power can be neglected in comparison to the accretion of phantom fluid onto the BH, particularly at later stage, we limit the analysis in the remaining sections to the interval $t \leq t_*$, far from the big rip singularity, and then purely classical treatment will be considered.

4 Thermodynamics with varying w

We now consider the thermodynamics properties of the solution described in section 2, with a variable EoS parameter, $p(a) = w(a)\rho(a)$. The particle fluid and entropy fluid currents, N^μ and S^μ are given by

$$N^\mu = nu^\mu, \quad S^\mu = su^\mu,$$

where n and s are the densities of particle number and entropy, respectively. The conservations laws, $T^{\mu\nu}_{;\nu} = 0$, $N^\mu_{;\mu} = 0$ and $S^\mu_{;\mu} = 0$, computed on the background given by (2), give the following set of differential equations

$$\dot{\rho} + \frac{\dot{R}_A}{R_A}\rho + \frac{3}{2}H(\rho + p) = 0, \quad (26)$$

$$\dot{n} + \left(\frac{\rho}{\rho + p} \frac{\dot{R}_A}{R_A} + \frac{3}{2}H \right) n = 0, \quad (27)$$

$$\dot{s} + \left(\frac{\rho}{\rho + p} \frac{\dot{R}_A}{R_A} + \frac{3}{2}H \right) s = 0. \quad (28)$$

The solutions of the above equations are

$$\rho(a) = \rho_0 \left[\frac{R_A(a_0)}{R_A(a)} \right] \left[\frac{a_0^{\frac{3}{2}(1+w_0)}}{a^{\frac{3}{2}(1+w(a))}} \right] \exp \left[\frac{3}{2} \int_{a_0}^a da w'(a) \ln a \right], \quad (29)$$

$$n(a) = n_0 \left(\frac{a_0}{a} \right)^{\frac{3}{2}} \exp \left[\int_a^{a_0} \frac{R'(a)}{(1+w(a))R(a)} da \right], \quad (30)$$

$$s(a) = s_0 \left(\frac{a_0}{a} \right)^{\frac{3}{2}} \exp \left[\int_a^{a_0} \frac{R'(a)}{(1+w(a))R(a)} da \right], \quad (31)$$

where the prime stands for derivative with respect to the scale factor, and ρ_0 , n_0 , s_0 are the present day values of the corresponding quantities assumed to be positive definite. Here we note the important corrections due to the presence of the black hole.

In the pure dark energy-dominated universe, $M_0 = 0$, we have $R_B = 0$ and $R_C = 1/H$. Consequently, using the equations of motion (11), we show that

$$\frac{R'_C(a)}{R_C(a)} = \frac{3}{2} (1 + w(a)) H(a). \quad (32)$$

Substituting in Eqs.(29-31) we obtain [33]

$$\rho(a) = \rho_0 \left[\frac{a_0^{3(1+w_0)}}{a^{3(1+w(a))}} \right] \exp \left[3 \int_{a_0}^a da w'(a) \ln a \right], \quad n(a) = n_0 \left(\frac{a_0}{a} \right)^3, \quad s(a) = s_0 \left(\frac{a_0}{a} \right)^3. \quad (33)$$

Now, assuming that $\rho = \rho(T, n)$, $p = p(T, n)$ and using the Gibbs law

$$T \left(\frac{\partial p}{\partial T} \right)_n = p + \rho - n \left(\frac{\partial \rho}{\partial n} \right)_T, \quad (34)$$

combined with

$$\dot{\rho} = \dot{n} \left(\frac{\partial \rho}{\partial n} \right)_T + \dot{T} \left(\frac{\partial \rho}{\partial T} \right)_n, \quad (35)$$

and the continuity equation, we obtain

$$\begin{aligned} & \left[\left(\frac{3}{2} H + \frac{1}{(1+w(a))} \frac{\dot{R}_A}{R_A} \right) T(a) w(a) + \dot{T}(a) \right] \left(\frac{\partial \rho}{\partial a} \right)_n \\ &= - \left(\frac{3}{2} H + \frac{1}{(1+w(a))} \frac{\dot{R}_A}{R_A} \right) T(a) \rho(a) w'(a). \end{aligned} \quad (36)$$

Calculating $\left(\frac{\partial \rho}{\partial a} \right)_n$ from Eq.(29), we get the equation governing the evolution of temperature

$$w'(a) T(a) - \left[\frac{3}{2a} (1 + w(a)) + \frac{R'_A}{R_A} \right] T(a) w(a) = (1 + w(a)) T'(a). \quad (37)$$

Solving for $w(a) \neq -1$, we obtain

$$T(a) = T_0 \left[\frac{w(a) + 1}{w_0 + 1} \right] \left[\frac{a_0^{3w_0/2}}{a^{3w/2}} \right] \exp \left[\int_{a_0}^a da \left[\frac{3}{2} w'(a) \ln a - \frac{w(a)}{(1+w(a))} \frac{R'_A(a)}{R_A(a)} \right] \right]. \quad (38)$$

Using this expression, we write the energy density as a function of temperature

$$\rho(a) = \rho_0 \left[\frac{T(a)}{T_0} \frac{(w_0 + 1)}{(w(a) + 1)} \right] \left(\frac{a_0}{a} \right)^{\frac{3}{2}} \exp \left[- \int_{a_0}^a \frac{da}{(1+w(a))} \frac{R'_A(a)}{R_A(a)} \right]. \quad (39)$$

Extracting the scale factor from (18), we obtain the generalized Stefan-Boltzmann law

$$\begin{aligned} \rho(T) = & \rho_0 \left[\frac{T(a)}{T_0} \frac{(w_0 + 1)}{(w(a) + 1)} \right]^{\frac{w(a)+1}{w(a)}} a_0^{\frac{3}{2} \frac{w(a)-w_0}{w(a)}} \exp \left[-\frac{3}{2w(a)} \int_{a_0}^a da w'(a) \ln a \right] \\ & \times \exp \left\{ \left[\frac{1}{w(a)} \int_{a_0}^a da \frac{w(a)}{(1 + w(a))} - \int_{a_0}^a \frac{da}{(1 + w(a))} \right] \frac{R'_A(a)}{R_A(a)} \right\}. \end{aligned} \quad (40)$$

Let us now reproduce the expressions of the temperature and energy density in the pure spatially flat FRW universe. Indeed, substituting (32) in (40) and (18), we obtain

$$\rho(T) = \rho_0 \left[\frac{T(a)}{T_0} \frac{(w_0 + 1)}{(w(a) + 1)} \right]^{\frac{w(a)+1}{w(a)}} a_0^{\frac{3}{2} \frac{w(a)-w_0}{w(a)}} \exp \left[\frac{3}{w(a)} \int_a^{a_0} da w'(a) \ln a \right] \quad (41)$$

and

$$T(a) = T_0 \left[\frac{w(a) + 1}{w_0 + 1} \right] \left[\frac{a_0^{3w_0}}{a^{3w}} \right] \exp \left[-3 \int_a^{a_0} da w'(a) \ln a \right], \quad (42)$$

which are exactly the relations obtained in [33].

Let us now scrutinize the behavior of the temperature when $w(a)$ crosses -1. In this case, Eq.(37) becomes

$$\left[w' + \frac{R'_A(a)}{R(a)} \right]_{\text{PDL}} T(a) = 0, \quad (43)$$

where $w'|_{\text{PDL}}$ is the value of $w(a)$ at the phantom divide line (PDL) and C is constant of integration. The solution of (43) is:

$$R_{\text{PDL}}(a) = C e^{-w'|_{\text{PDL}}(a-a_0)}, T(a) \neq 0, \quad \text{or} \quad T(a) = 0. \quad (44)$$

Here, we note that unlike the standard vacuum solution with $T = 0$ [33], the solution considered in this paper allows for a vacuum solution with a non-zero temperature. This was expected, since this temperature is associated with the AH of the black hole in the absence of dark field. The zero temperature vacuum state is recovered by setting $M_0 = 0$. On the other it is important to observe that the expressions of temperature and energy density are regular everywhere including the phantom divide crossing. This can be easily verified by substituting the solution of the AH at the PDL, $R(a)_{\text{PDL}} = C e^{-w'|_{\text{PDL}}(a-a_0)}$ in Eqs.(18,39)

$$T_{\text{PDL}}(a) = T_0 a_0^{3w_0/2} a^{3/2} \exp \left[-1 - w_0 + \frac{3}{2} w'|_{\text{PDL}} \int_{a_0}^a da \ln a \right], \quad (45)$$

and

$$\rho_{\text{PDL}}(a) = \rho_0 a_0^{3(w_0+1)/2} \exp \left[-1 - w_0 + \frac{3}{2} w'|_{\text{PDL}} \int_{a_0}^a da \ln a \right]. \quad (46)$$

Obviously, Eq.(45) shows that the temperature remains positive when $w(a) \rightarrow -1^\pm$. Now, making the following ansatz for the AH

$$R_A(a) = C e^{-w'|_{\text{PDL}}(a-a_0)} f_A(a), \quad (47)$$

with $f_A(a) \xrightarrow{w \rightarrow -1^\pm} 1$, we rewrite the temperature as

$$T(a) = T_{\text{PDL}} \left[\frac{w(a) + 1}{[w(a) + 1]_{\text{PDL}}} \right] a^{-\frac{3}{2}(w(a)+1)} \times \exp \left[-\frac{3}{2} w'(a)|_{\text{PDL}} \int_{a_0}^a da \ln a + \int_{a_0}^a da \left[\frac{3}{2} w'(a) \ln a - \frac{w(a)}{(1+w(a))} \frac{f'_A(a)}{f_A(a)} \right] \right], \quad (48)$$

which shows that the temperature is always positive definite regardless the value of $w(a)$. When $M_0 = 0$ the temperature is positive for $w(a) > 1$, negative for $w(a) < -1$ and zero for $w(a) = -1$ [33].

Now, we calculate the chemical potential defined by the Euler relation [34]

$$\mu(a) = \frac{(w(a) + 1) \rho(a)}{n(a)} - \frac{T(a) s(a)}{n(a)}. \quad (49)$$

Using the expressions of $\rho(a)$, $n(a)$, $s(a)$ and $T(a)$ we obtain

$$\mu(a) = \mu_0 \left[\frac{w(a) + 1}{w_0 + 1} \right] \left[\frac{a_0^{3w_0/2}}{a^{3w(a)/2}} \right] \exp \left\{ \int_{a_0}^a da \left[\frac{3}{2} w'(a) \ln a - \frac{w(a)}{(1+w(a))} \frac{R'_A(a)}{R_A(a)} \right] \right\}, \quad (50)$$

where

$$\mu_0 = \frac{(w_0 + 1) \rho_0}{n_0} - \frac{T_0 s_0}{n_0} \quad (51)$$

is the present day chemical potential. We note that in general the sign of μ_0 can be arbitrary, and consequently the sign of $\mu(a)$.

The entropy of the universe can be derived from (49) and is given by

$$s(a) = \frac{(w(a) + 1) \rho(a) - \mu(a) n(a)}{T(a)}. \quad (52)$$

Using again the relations for $\rho(a)$, $T(a)$, $\mu(a)$ and $n(a)$ we obtain

$$s(a) = s_0 \frac{a_0^{\frac{3}{2}}}{a^{\frac{3}{2}}} \exp \left[\int_a^{a_0} \frac{R'_A(a)}{(1+w(a)) R_A(a)} da \right], \quad (53)$$

where the present day entropy is

$$s_0 = \left[\frac{(w_0 + 1) \rho_0}{T_0} - \frac{\mu_0 n_0}{T_0} \right]. \quad (54)$$

Now, defining the comoving volume by

$$V(a) = a^{\frac{3}{2}} \exp \left[\frac{1}{2} \int^a \frac{R'_A(a)}{(1 + w(a)) R_A(a)} da \right], \quad (55)$$

we derive from Eq.(53) the usual entropy conservation law

$$s(a)V(a) = s_0 V_0. \quad (56)$$

As we did for the energy density, we write the entropy in terms of temperature as

$$\begin{aligned} s(T) = s_0 & \left[\frac{T(a)}{T_0} \frac{w_0 + 1}{w(a) + 1} \right]^{\frac{1}{w(a)}} a_0^{\frac{3}{2} \left(\frac{w(a) - w_0}{w(a)} \right)} \exp \left[-\frac{3}{2w(a)} \int_{a_0}^a da w'(a) \ln a \right] \\ & \times \exp \left[\left[\frac{1}{w(a)} \int_{a_0}^a da \frac{w(a)}{(1 + w(a))} - \int_{a_0}^a \frac{da}{(1 + w(a))} \right] \frac{R'_A(a)}{R_A(a)} \right]. \end{aligned} \quad (57)$$

Let us now consider the expressions of the chemical potential and entropy at the phantom divide crossing. Using (3.21) we easily show that

$$\mu_{\text{PDL}}(a) = \mu_0 a_0^{3w_0/2} a^{3/2} \exp \left\{ -1 - w_0 + \frac{3}{2} w'|_{\text{PDL}} \int_{a_0}^a da \ln a \right\}, \quad (58)$$

and

$$s_{\text{PDL}}(a) = s_0 \frac{a_0^{\frac{3}{2}}}{a^{\frac{3}{2}}} \exp \left\{ -1 - w_0 + \frac{3}{2} w'|_{\text{PDL}} \int_{a_0}^a da \ln a \right\}. \quad (59)$$

Note that using relations (45), (58) and (59) we easily verify that the Euler relation remains valid at the phantom divide line.

Finally, let us list the corresponding relations for the energy density, temperature, entropy and chemical potential in the case $w(a) = w = \text{const}$,

$$T(a) = T_0 \left[\frac{a_0}{a} \right]^{3w/2} \left[\frac{R_A(a_0)}{R_A(a)} \right]^{\frac{w}{1+w}}, \quad (60)$$

$$\rho(a) = \rho_0 \left[\frac{a_0}{a} \right]^{\frac{3}{2}(1+w)} \left[\frac{R_A(a_0)}{R_A(a)} \right], \quad (61)$$

$$\mu(a) = \mu_0 \left[\frac{a_0}{a} \right]^{3w/2} \left[\frac{R_A(a_0)}{R_A(a)} \right]^{\frac{w}{1+w}}, \quad (62)$$

$$s(a) = s_0 \left[\frac{a_0}{a} \right]^{\frac{3}{2}} \left[\frac{R_A(a_0)}{R_A(a)} \right]^{\frac{1}{1+w}}, \quad (63)$$

$$\rho(T) = \rho_0 \left[\frac{T(a)}{T_0} \right]^{\frac{1+w}{w}}, \quad (64)$$

$$\mu(T) = \mu_0 \left(\frac{T(a)}{T_0} \right), \quad (65)$$

$$s(T) = s_0 \left[\frac{T(a)}{T_0} \right]^{\frac{1}{w}} = s_0 \left[\frac{\rho}{\rho_0} \right]^{\frac{1}{1+w}}. \quad (66)$$

It is interesting to note that the dependence of the thermodynamics parameters on the temperature is of the same form as in standard thermodynamics of purely expanding FRW universe.

Before ending this section, let us reconsider explicitly the avoidance of the big rip singularity in a phantom dominated universe. Using Eqs.(26,61) we obtain

$$a(t) = a_0 \left[\frac{3}{2} H_0 (1+w) (t - t_s) \right]^{\frac{2}{3(1+w)}}, \quad (67)$$

$$H(t) = \frac{2}{3(1+w)(t - t_s)}, \quad (68)$$

where the big rip time is

$$t_s = t_0 - \frac{2}{3H_0(1+w)}. \quad (69)$$

Using the constraint (13) on $H(t)$, in phantom dominated era, one finds

$$(t_s - t) \geq \frac{16Gm_H(t)}{3(|w| - 1)}. \quad (70)$$

Then using the FRW equations and (15), we find

$$\rho < \frac{(t - t_s)^{-1}}{8\pi G^2 m_H (1+w)}, \quad |p| < \frac{w(t_s - t)^{-1}}{8\pi G^2 m_H (1+w)}. \quad (71)$$

Substituting (70) in these relations we reproduce the relations (16). Hence, the universe evolves towards a state where a , ρ , p and higher derivatives are finite, without ever reaching the big rip singularity. This final state is reached in a finite time given by

$$t_* = t_s + \frac{2 \left[8GH_0 M_0 a_0^{\frac{1}{2}(5+3w)} \right]^{\frac{1+3w}{3(1+w)}}}{3H_0(1+w)}. \quad (72)$$

This behavior of the universe is similar to that observed in the framework of generalized uncertainty principle (GUP) corrected FRW universe [35].

5 Stability of the solution

In the following, we focus on the regime of cosmic dynamics where the universe undergoes a phase of quasi-exponential expansion, such that $\dot{H}/H^2 \ll 1$, and examine the stability of the solutions obtained above at the crossing of the phantom divide line when quantum effects due to conformal anomaly are taken into account. In general, the conformal anomaly is given by [36]

$$T = b \left(F + \frac{2}{3} \square R \right) + b' G + b'' \square R, \quad (73)$$

where $F = C_{\mu\nu\lambda\kappa} C^{\mu\nu\lambda\kappa} = \frac{1}{3} R^2 - 2R_{\mu\nu} R^{\mu\nu} + R_{\mu\nu\lambda\kappa} R^{\mu\nu\lambda\kappa}$ is the square of the Weyl tensor and G the Gauss-Bonnet invariant, $G = R^2 - 4R_{\mu\nu} R^{\mu\nu} + R_{\mu\nu\lambda\kappa} R^{\mu\nu\lambda\kappa}$, and the coefficient b and b' are

$$b = \frac{1}{120 (4\pi)^2} (n_0 + 6n_{1/2} + 12n_1), \quad (74)$$

$$b' = \frac{1}{360 (4\pi)^2} (n_0 + 11n_{1/2} + 62n_1), \quad (75)$$

where n_0 , $n_{1/2}$ and n_1 are the number of scalar, Dirac fermion and vector fields, respectively.

Since we have the condition (13), we just give the expressions of $\square R$, F and G to leading order in $Gm_H H$ as

$$F \sim O((Gm_H H)^2), \quad \square R \sim O((Gm_H H)^2), \quad (76)$$

$$G \simeq 42b' H^4 + 128b' H^5 Gm_H. \quad (77)$$

Now assuming that

$$T_a = -\rho_a + 3p_a. \quad (78)$$

and using the conservation law

$$\dot{\rho}_a + 3H(\rho_a + p_a) = 0, \quad (79)$$

one finds

$$4\rho_a + \frac{1}{H} \frac{d\rho_a}{dt} = -T_a. \quad (80)$$

Then ρ_a and p_a are given by

$$\rho_a = -\frac{1}{a^4} \int dt a^4 H T_a, \quad p_a = \frac{T_a + \rho_a}{3}. \quad (81)$$

In the quasi-exponential expansion we can write $\rho_a \sim p_a \sim T_a$, and then we have

$$\rho_a \sim p_a \sim 42b' H^4 + 128b' H^5 Gm_H. \quad (82)$$

On the other hand the fluid density energy can be approximated by

$$\rho_f \sim p_f \sim \frac{3H^2}{8\pi G} + \frac{3m_H H^3}{4\pi G}. \quad (83)$$

Taking the present day values of the Hubble parameter, $H_0 \sim 10^{-33}$ eV, and $G \sim 10^{-56} (\text{eV})^{-2}$, it is easy to verify that the quantum correction are very small when crossing the phantom divide line. Even we consider BH masses at the end of the expansion of the order of $\sim 10^{23} M_\odot$, the correction term to the standard result is insignificant. Hence, the solutions given in section 4 are stable under the conformal anomaly quantum corrections.

6 Accretion of phantom fluid and constraints on the GSL

We now consider the problem of the validity of the GSL when the effect of the back-reaction effect of the phantom fluid on the black hole is taken into account. We restrict our study to the scenario of a black hole with a small quasi-local mass immersed in phantom fluid dominated era. In this case only the cosmological AH contributes, and the total entropy consists essentially of the sum of entropy of the cosmological AH and the entropy of the phantom fluid in thermal equilibrium with the cosmological AH. Indeed, using (66) we have

$$S = \left[\frac{\pi R_C^2}{G} + s_0 V \left(\frac{\rho}{\rho_0} \right)^{\frac{1}{1+w}} \right]. \quad (84)$$

The first term is the entropy of the cosmological AH and the second term is the phantom fluid entropy inside a comoving volume V . Taking the derivative with respect to time and using the following approximation

$$R_C \approx \frac{1}{H} - \frac{2G\dot{m}_h}{H}, \quad (85)$$

along with the relations $\dot{m}_h = m_h \ddot{a}/a$, $\ddot{a} = a(H^2 + \dot{H})$, we obtain to leading order in the Hawking-Hayward quasi-local mass

$$\begin{aligned} \dot{S} = & -\frac{2\pi}{GH} \left[\frac{\dot{H}}{H^2} + 2G\dot{m}_H \left(1 - \frac{\dot{H}}{H^2} \right) \right] \\ & + \frac{s_0 V H}{(1+w)} \left(\frac{T}{T_0} \right)^{\frac{1}{w}} \left[\frac{\dot{H}}{H^2} + 2G\dot{m}_H \left(1 + \frac{\dot{H}}{H^2} \right) - \frac{3}{2}(1+w) \right], \end{aligned} \quad (86)$$

where we have used the continuity equation (26) and (66). In order to satisfy the GSL, $\dot{S} \geq 0$, the quantity inside the square brackets must be positive definite, yielding to the condition

$$2\dot{m}_H \geq \frac{\frac{2\pi\dot{H}}{GH^2} - \frac{s_0 V H^2}{(1+w)} \left(\frac{\rho}{\rho_0}\right)^{\frac{1}{1+w}} \left[\frac{\dot{H}}{H^2} - \frac{3}{2}(1+w)\right]}{\frac{s_0 V H^2}{(1+w)} \left(\frac{\rho}{\rho_0}\right)^{\frac{1}{1+w}} G \left(1 + \frac{\dot{H}}{H^2}\right) - 2\pi \left(1 - \frac{\dot{H}}{H^2}\right)}. \quad (87)$$

Now substituting $\frac{\dot{H}}{H^2} = -\frac{3}{2}(1+w)$, we rewrite the constraint on the GSL as

$$G\dot{m}_H \geq -3 \frac{(1+w)}{(1+3w)} \left[\frac{\mathcal{A} - \frac{1+w}{2}}{\mathcal{A} + \frac{(5+3w)(1+w)}{1+3w}} \right], \quad (88)$$

with $\mathcal{A} = \frac{s_0 V H^2 G}{2\pi} \left(\frac{T}{T_0}\right)^{\frac{1}{w}}$.

We know from Eq.(12) that the Hawking-Hayward quasi-local mass is an increasing function of time in an expanding universe, $\dot{m}_H \geq 0$. Then, imposing the positiveness of the RHS of Eq.(88) and assuming the positiveness of the entropy, we have one solution given by

$$-\frac{(5+3w)(1+w)}{(1+3w)} > \mathcal{A} \geq 0, \quad (89)$$

which is satisfied for an EoS parameter in the range $w \leq -5/3$. Using the expression of s_0 given in (54) and defining $\alpha_0 = -\mu_0 n_0 / \rho_0$, an immediate consequence of the positiveness of the entropy is that $w \geq -1 - \alpha_0$. Since $w \leq -5/3$, we get a lower bound estimate to the present day value of the chemical potential,

$$\alpha_0 \geq \frac{2}{3}, \quad (90)$$

which show that the chemical potential in phantom energy dominated-era is strictly negative. Now, with $w = -2$ and the present day parameters, we obtain the following bound on the entropy

$$s_0 < 3.5 \times 10^{-4} \text{ GeV}^{-3}. \quad (91)$$

Or in terms of α_0 and using (90), we have

$$\frac{2}{3} \leq \alpha_0 < 1 + 1.26 \times 10^{43} T_0. \quad (92)$$

Now, if we take $T_0 \simeq 10^{-19} \text{ GeV}$ we have $\alpha_0 \lesssim 10^{24}$. Here we would like to point that the bounds on the present day values of the entropy and the parameter α_0 are of the same order of that obtained by Lima et al. [37]. However, the calculation in this paper is performed by ignoring the back-reaction of the phantom fluid on the black hole. In fact, they used the Schwarzschild metric, which cannot describe the properties of black holes embedded in an expanding FRW universe. As we explicitly show, taking into account the back-reaction effect, even in a low density background, leads to a drastic constraint on the EoS parameter and rule out the approach based on zero chemical potential [18].

Now, using the obvious relation $m_H = \dot{m}_H/H$, we obtain the quasi-local black hole mass above which the accretion of phantom fluid onto the black hole is permitted

$$m_H \geq -\frac{3}{GH} \frac{(1+w)}{(1+3w)} \left[\frac{\mathcal{A} - \frac{1+w}{2}}{\mathcal{A} + \frac{(5+3w)(1+w)}{1+3w}} \right]. \quad (93)$$

Using the constraint on the GSL (89), we obtain

$$m_H \gtrsim m_{H,crit} = -\frac{6}{GH} \left[\frac{\mathcal{A} - \frac{1+w}{2}}{(5+3w)} \right]. \quad (94)$$

In terms of α_0 and the present day quantities, the critical quasi-local mass is then

$$m_{H,crit} = -42.27 \times 10^{-20} \left[\frac{1+w+\alpha_0}{5+3w} \right] \frac{M_\odot}{T_0}. \quad (95)$$

For large values of $\alpha_0 \sim 10^{24}$ and $T_0 \approx 10^{-19}$ GeV, the critical quasi-local mass is approximately $m_{H,crit} \approx 10^{23} M_\odot$. This is a huge value, allowing all black holes in the universe to accrete phantom fluid. We note that in section 3, we have shown that $m_{H,crit}$ is the maximal allowed mass which prevents the engulfing of the universe by the BH. On the other hand small values of α_0 give critical BH mass of the order of the solar mass. For instance, taking $\alpha_0 = 2$, $w = -2$ and $T_0 \approx 10^{-19}$ GeV, we obtain $m_{H,crit} \approx 4.2 M_\odot$. Our results are similar to that obtained in [37], where the EoS parameter is restricted to values less than -1 and the back-reaction effect ignored. We have plotted the present day critical quasi-local mass with respect to w for different values of the parameter α_0 . Obviously, the value of the parameter α_0 is very important in determining a critical quasi-local mass of the order of the solar mass. Finally, if we set the chemical potential to zero, the critical mass reduces to

$$m_{H,crit} = -\frac{3}{2\pi} \frac{(1+w)}{(5+3w)} \frac{\rho_0 V H}{T_0} \left(\frac{T}{T_0} \right)^{\frac{1}{w}}, \quad (96)$$

which is negative for $w < -5/3$. This result is expected from the relation (90).




Figure 3: Variation of the critical Hawking-Hayward quasi-local mass with respect to w for different values of the parameter α_0 .

7 Conclusion

In summary, we have investigated the thermodynamical properties of black holes immersed in an expanding spatially flat FRW universe, for general EoS parameter $w(a)$. Particularly, we found that the temperature of the dark fluid is always positive regardless the value of the time varying EoS parameter, and that the instantaneous vacuum state is characterized by non-zero temperature, entropy and chemical potential, respectively. An other important result is that all the thermodynamics parameters are regular at the phantom divide crossing. We have also analyzed the accretion process of phantom fluid onto black holes with small Hawking-Hayward quasi-local mass, and the constraints on the validity of the GSL. Particularly, assuming the positiveness of the phantom fluid entropy, we have shown that phantom fluid may have a negative definite chemical potential in order to satisfy the GSL. We have also obtained, for an EoS parameter within the interval $w < -5/3$, a critical quasi-local mass of the black hole, above which the GSL is always protected. The present analysis, show that taking into account the back-reaction effect of the phantom fluid on the black hole, even in a low density background, leads naturally to positive temperature and negative chemical potential, and may contributes to resolve the controversy on the subject [18, 37].

Acknowledgments

The author was supported by the Algerian Ministry of High Education and Scientific Research under the CNEPRU project N°: D01720070033.

References

- [1] S. Perlmutter et al. [*Supernovae Cosmology Project Collaboration*], *Astrophys. J.* **517**, 565 (1999) [arXiv:astro-ph/9812133]
- [2] A. G. Riess et al. [Supernovae Search Team Collaboration], *Astron. J.* **116**, 1009 (1998) [arXiv:astro-ph/9805201].
- [3] S. Capozziello, *Int. J. Mod. Phys. D* **11**, 483 (2002); S. Capozziello, V.F. Cardone, S. Carloni and A. Troisi, *Int. J. Mod. Phys. D* **12**, 1969 (2003); S. Capozziello, V.F. Cardone and A. Troisi, *Phys. Rev. D* **71**, 043503 (2005); S. Carloni, P.K.S. Dunsby, S. Capozziello and A. Troisi, *Class. Quant. Grav.* **22**, 4839 (2005); S. Nojiri and S.D. Odintsov, *Phys. Lett. B* **576**, 5 (2003); S. Nojiri and S.D. Odintsov, *Mod. Phys. Lett. A* **19**, 627 (2003); S. Nojiri and S.D. Odintsov, *Phys. Rev. D* **68**, 12352 (2003); S.M. Carroll, V. Duvvuri, M. Trodden and M. Turner, *Phys. Rev. D* **70**, 043528 (2004); G. Allemandi, A. Borowiec and M. Francaviglia, *Phys. Rev. D* **70**, 103503 (2004).
- [4] G. Dvali, G. Gabadadze and M. Porrati, *Phys. Lett. B* **485**, 208 (2000); G. Dvali and G. Gabadadze, *Phys. Rev. D* **63**, 065007 (2001).
- [5] R. R. Caldwell, M. Kamionkowski and N. N. Weinberg, *Phys. Rev. Lett.* **91**, 071301 (2003) [arXiv:astro-ph/0302506].
- [6] B. McInnes, *JHEP* **0208**, 029 (2002) [arXiv:hep-th/0112066].
- [7] E. Elizalde, S. Nojiri and S. D. Odintsov, *Phys. Rev. D* **70**, 043539 (2004) [arXiv:hep-th/0405034];
- [8] S. Nojiri and S. D. Odintsov, *Phys. Lett. B* **595**, 1 (2004) [arXiv:hep-th/0405078].
- [9] S. Nojiri and S. D. Odintsov, *Phys. Rev. D* **70**, 103522 (2004) [arXiv:hep-th/0408170].
- [10] S. Nojiri and S. D. Odintsov and S. Tsujikawa, *Phys. Rev. D* **70**, 043520 (2004) [arXiv:hep-th/0401025];
- [11] J. M. Cline, S. Y. Jeon and G. D. Moore, *Phys. Rev. D* **70**, 043543 (2004) [arXiv:hep-ph/0311312]; P. Singh, M. Sami and N. Dadhlich, *Phys. Rev. D* **68**, 0.23522 (2003) [arXiv:hep-th/030510].
- [12] S. Nojiri and S. D. Odintsov, *Phys. Lett. B* **565**, 1 (2003) [arXiv:hep-th/0304131]; S. Nojiri and S. D. Odintsov, *Phys. Lett. B* **562**, 147 (2003) [arXiv:hep-th/0303117]; A. E. Schulz and M. J. White, *Phys. Rev. D* **64**, 043514 (2001) [arXiv:astro-ph/0104112].

- [13] A. Melchiorri, I. Mersini-Houghton, C. J. Odman and M. Trodden, Phys. Rev. D **68**, 043509 (2003) [arXiv:astro-ph/0211522].
- [14] R. Brustein, Phys. Rev. Lett. **84**, 2072 (2000) [arXiv:gr-qc/9904061]; M. Li, Phys. Lett. B **603**, 1 (2004); P. C. W. Davies, Class. Quant. Grav. **4**, L225 (1987); Class. Quant. Grav. **5**, 1349 (1988); R. Bousso, Phys. Rev. D **71**, 064024 (2005); S. Nojiri and S. D. Odintsov, Phys. Rev. D **72**, 063505 (2005); Q. Huang and M. Li, JCAP **0408**, 013 (2004); H. Mohseni-Sadjadi and M. Honardoost, Phys. Lett. B **647**, 231 (2007) [arXiv:gr-qc/0609079]; P. F. Gonzalez-Diaz and C. L. Siguenza, Nucl. Phys. B **697**, 363 (2004) [arXiv:astro-ph/0407421].
- [15] I. Brevik, S. Nojiri, S.D. Odintsov and L. Vanzo, Phys. Rev. D **70**, 043520 (2004) [arXiv:hep-th/0401073]; J. A. S. Lima and J. S. Alcaniz, Phys. Lett. B **600**, 191 (2004) [arXiv:astro-ph/0402265].
- [16] G. Izquierdo and D. Pavljuchenko, Phys. Lett. B **633**, 420 (2006) [arXiv:astro-ph/0505601]; Phys. Lett. B **639**, 1 (2006) [arXiv:gr-qc/0606014].
- [17] J. A. S. Lima and S. H. Pereira, Phys. Rev. D **78**:083504, 2008 [arXiv:0801.0323]; S. H. Pereira and J. A. S. Lima, Phys. Lett. B **669**, 266 (2008). [arXiv:0806.0682]; S. H. Pereira, [arXiv:0806.3701].
- [18] J. A. de Freitas Pacheco and J. E. Horvath, Class. Quant. Grav. **24**, 5427 (2007) [arXiv:0709.1240];
- [19] E. Babichev, V. Dokuchaev, and Yu. Eroshenko, Phys. Rev. Lett. **93**, 021102 (2004) [arXiv:gr-qc/0402089].
- [20] V. Faraoni and A. Jaques, Phys. Rev. D **76**, 063510 (2007) [arXiv:gr-qc/0707135v1].
- [21] C. Gao, X. Chen, V. Faraoni and Y-G. Shen, Phys. Rev. D **78**, 024008 (2008) [arXiv:0802.1298].
- [22] G. C. McVittie, Mon. Not. R. Astr. Soc. **93**, 325 (1933).
- [23] R. Sussman, Gen. Rel. Grav. **17**, 251 (1985); M. Ferraris, M. Francaviglia and A. Spallucci, Nuovo Cimento B **111**, 1031 (1996);
- [24] B. C. Nolan, Class. Quantum. Grav. **16**, 1227 (1999); B. C. Nolan, Class. Quantum. Grav. **16**, 3183 (1999) [arXiv:gr-qc/9907018]; B. C. Nolan, Phys. Rev. D **58**, 064006 (1998).
- [25] P. Martin-Moruno, J. A. Jimenez Madrid and P. F. Gonzalez-Diaz [arXiv:astro-ph/0603761v1].

- [26] M. J. Rees and M. Volonteri, *Massive Black Holes: formation and evolution in “Black Holes: from Stars to Galaxies – across the Range of Masses”, Proceedings IAU Symposium No. 238, V. Karas & G. Matt, eds. (2006).*
- [27] S. A. Hayward, Phys. Rev. D **70**, 104027 (2004) [arXiv:gr-qc/0408008]; Phys. Rev. Lett. **93**, 251101 (2004) [arXiv:gr-qc/0404077]; Phys. Rev. Lett. **81**, 4557 (1998) [arXiv:gr-qc/9807003]; Phys. Rev. D **53**, 1938 (1996); Class. Quant. Grav. **11**, 3025 (1994); Phys. Rev. D **49**, 6467 (1994); [arXiv:gr-qc/9303006]; S. A. Hayward, S. Mukohyama and M. C. Ashworth, Phys. Lett. A **256**, 347 (1999) [arXiv:gr-qc/9810006]; Mukohyama and S. A. Hayward, Class. Quant. Grav. **17**, 2153 (2000).
- [28] I. Arraut, D. Batic, and M. Nowakowski, Class. Quant. Grav. **26**, 125006 (2009) [arXiv:0810.5156].
- [29] J. D. Bekenstein, Phys. Rev. D **23**, 287 (1981); C. Massa, Am. J. Phys. **54**, 733 (1986); C. Massa, Am. J. Phys. **57**, 91 (1989).
- [30] A. Bonanno and M. Reuter, Phys. Rev. D **62**, 043008 (2000) [arXiv:hep-th/0002196].
- [31] V. Faraoni, Phys. Rev. D **80**, 044013 (2009) [arXiv:0907.4473].
- [32] Cheng-Yi Sun, [arXiv:0906.3783v5].
- [33] Emmanuel N. Saridakis, Pedro F. Gonzalez-Diaz and Carmen L. Siguenza, Class. Quant. Grav. **26**, 165003 (2009) [arXiv:0901.1213].
- [34] H. B. Callen, *Thermodynamics and an Introduction to Thermostatistics*, 2nd edition, Wiley, New-York (1985).
- [35] K. Nouicer, *Avoidance of the big rip singularity in GUP-corrected Friedmann-Robertson-Walker spacetime*, in progress.
- [36] M. J. Duff, Class. Quant. Grav. **11**, 1387 (1994) [arXiv:hep-th/9308075]; K. Bamba, C.-Q. Geng, S. Nojiri and S. D. Odintsov, Phys. Rev. D **79**, 083014 (2009) [arXiv:0810.4296].
- [37] J. A. S. Lima, S. H. Pereira, J. E. Horvath and D. C. Guariento, [arXiv:0808.0860].